Reasons for requiring new design-levels for Christchurch region

- Greatly increased seismicity rates compared to low pre-September 2010 rates
 - sequence likely to continue at reducing rates for years rather than months
- Stronger ground motions in CBD than given by models
- Possible causes
 - stress-drops
 - directivity
 - site effects (long-period peak)

New interim seismic hazard model for Christchurch

- Accounts for ongoing earthquake sequence
 - includes decreasing rates with time
- Shallow depth
 - 5 km rather than 10 km minimum in model
- Increases ground-motions
 - uses increased stress-drop, possibly surrogate for other effects such as directivity for which suitable models not readily available
- Adjusts for dominance of magnitude 5-6 events
 - advice from Engineering Advisory Group that magnitudes less than 5.5 of little significance for engineered structures
- Limited to 1.5s period maximum until long-period modelling of site effects available

Royal Commission 18 October 2011

GNS Science and University of Canterbury

Maximum magnitudes used for distributed seismicity in hazard calculations

- Seismicity model consists of fault sources (from geology) and a grid of distributed seismicity (from historical earthquake catalogue)
- Maximum magnitude of 7.0 in the Christchurch area for grid sources in 2000/2002 National Seismic Hazard Model (NSHM) used to develop NZS1170
- Larger magnitudes were associated with specific known faults e.g. Porters-Grey M7.5, Alpine M8.1
- Maximum magnitude for distributed seismicity increased to 7.2 in 2010 update of NSHM
- Post-February time-varying model for Canterbury has magnitudes up to 7.9 for distributed seismicity


Minimum allowable Z-factor = 0.13 from magnitude 6.5 earthquake at 20 km distance

- Z-factor usually corresponds to a spectral acceleration with an annual exceedance probability of 1/500
- Minimum allowable value Z=0.13 based on scenario earthquake motions
- Corresponds to 2/3 the 84th-percentile (one standard deviation above the median) motions for a magnitude 6.5 earthquake at 20 km distance
- Derived from survivability level in earthquakes that may not be apparent from geology in lowest seismicity regions
- 2/3 factor to convert from survivability to design-level Ultimate Limit State motions
- Comes into play only in lowest seismicity regions
 - north of Bombay Hills, and in south-eastern Otago
 - Christchurch Z=0.22 well above Z=0.13

Preliminary results Z=0.3

- Engineering Advisory Group requested appropriate Z-factor (sets level of design spectrum) to use with NZS1170 Deep Soil spectral shape (site conditions appropriate for most of Christchurch CBD)
- Usual normalisation is in terms of Shallow Soil spectral values
- Initial results gave Z=0.3 using average seismicity rate over next 50 years
 - 50 years is default design life for most structures
 - Z=0.22 in NZS1170
 - approximately 0.4 (or greater) for CBD motions in February

Initial hazard spectrum and Z=0.3 Deep Soil spectrum




Intended for engineered structures 0.5s-1.5s period 1/500 average annual exceedance probability (AEP) over next 50 years

Excludes long-period site effect around 2.5-4s

Latest model

- Latest model raises Z=0.30 to range Z=0.34-0.41
 - resetting of seismic activity following magnitude 6.0 aftershock on 13 June 2011
 - adjustment of some parameters from US to NZ defaults in new software (NZ uses larger rather than geometric mean of 2 horizontal components)
- Wellington Z=0.4
- February CBD motions approximately Z=0.4 or greater

1/500 AEP hazard spectrum over next 50 years and NZS1170 deep soil Z=0.34 and 0.41 spectra



Z=0.34 is lower bound for period range 0.5s-1.5s (Zmin)


Royal Commission 18 October 2011

GNS Science and University of Canterbury

Time-variation of Z-factor

- Decreasing seismicity causes Z-factor to decrease with time
- Current value greater than Z=0.34 average over next 50-years
- Later falls below Z=0.34 average

Variation with Z-factor over time



Z=0.34 average for next 50 yrs
Z=0.48 for 12 months
from 1 October 2011
Drops to 0.34 in year 8 (2019)
Z=0.3 beyond year 20
(close after year 15)

GNS Science and University of Canterbury